Research in Astron. Astrophys. **20116** Vol. **?** No. **XX**, 000–000
http://www.raa-journal.org *http://www.iop.org/journals/raa*

***R**esearch in*
***A**stronomy and*
***A**strophysics*

arXiv:1612.02993v1 [astro-ph.SR] 9 Dec 2016

Chromospheric activity on late-type star of V1355 Ori using Lijiang 1.8-m and 2.4-m telescopes

Qingfeng Pi[1,2,3], Liyun Zhang[1,2], Liang Chang[2,4], Xianming L. Han[5], Hongpeng Lu[1,2], XiLiang Zhang[2,4] and Daimei Wang[1,2]

[1] College of Physics/Department of Physics and Astronomy, Guizhou University, Guiyang 550025, P.R. China *piqingfeng@126.com; liy_zhang@hotmail.com*
[2] Key Laboratory for the Structure and Evolution of Celestial Objects, Chinese Academy of Sciences, Kunming 650011 P.R. China
[3] MingDe College, Guizhou University, Guiyang 550025, P.R. China
[4] Yunnan Observatories, Chinese Academy of Sciences, Kunming, 650011, P.R. China
[5] Dept. of Physics and Astronomy, Butler University, Indianapolis, IN 46208, USA

Received ?; accepted ?

Abstract We obtained new high-resolution spectra using the Lijiang 1.8-m and 2.4-m telescopes to discuss the chromospheric activities of V1355 Ori as indicated in the behaviour in the Ca II H&K, H_δ, H_γ, H_β, Na I D_1 D_2, H_α and Ca II infrared triplet (IRT) lines. The observed spectra show obvious emissions above continuum in the Ca II H&K lines, absorptions for the H_δ, H_γ, H_β, and Na I D_1, D_2 lines, variable behaviour (filled-in absorption, partial emission with a core absorption component, or emission above continuum) for the H_α line, and weak self-reversal emissions in the strong filled-in absorptions of the Ca II IRT lines. We used a spectral subtraction technique to analyse our data. The results show no excess emission in the H_δ and H_γ lines, very weak excess emissions for the Na I D_1, D_2 lines, excess emission in the H_β line, clear excess emission for the H_α line, and excess emissions for the Ca II IRT lines. The value of the ratio of EW_{8542}/EW_{8498} is in the range of 0.9 to 1.7, and it implies that the chromospheric activity might have been caused by plage events. The value of the ratio E_{H_α}/E_{H_β} is above 3, and it implies that the Balmer lines would arise from prominence-like material. We also found time variations of light curves of equivalent widths (EWs) of the chromospheric activity lines in the Na I D_1 D_2, Ca II IRT, and H_α lines in particular. These phenomena can be explained by plage events, which are consistent with the behaviour of chromospheric active indicators.

Key words: stars: chromosphere – stars: activity – stars: individual V1355 Ori

1 INTRODUCTION

V1355 Ori (BD -00 1147, HD 291095, K0-2 IV, 3.82 days) is a single-line spectroscopic RS CVn binary (Strassmeier et al. 1999; Strassmeier 2000; Strassmeier et al. 2000; Savanov & Strassmeier 2008; Eker et al. 2008; etc). It exhibits photospheric and chromospheric magnetic activities, which has been discovered at optical (Cutispoto et al. 1995; Strassmeier 2000; Savanov & Strassmeier 2008; etc), ultraviolet (Pounds et al. 1993), and X-ray wavelengths (Voges et al. 1999).

Table 1 Observation log of V1355 Ori using 1.8-m and 2.4-m telescopes.

Date	HJD(245,)	Exp. time	S/N										
	days	(s)	Ca II IRTλ8662	Ca II IRTλ8542	Ca II IRTλ8498	H_α	Metal line	Na I	H_β (4861 Å)	H_γ (4341 Å)	H_δ (4102Å)	Ca II H (3968Å)	Ca II K (3933Å)
20150122	7045.22327	3600	58	62	52	52	81	70					
20150124	7046.23188	1800	47	51	44	44	53	51					
20150127	7050.17989	2400	45	50	43	43	54	52					
20150204	7058.14409	3600	68	73	64	64	77	73					
20150206	7060.11388	3600	67	71	62	62	81	75					
20150209	7063.11839	1800	45	49	43	43	52	47					
20150210	7064.13065	2400	49	53	45	45	58	55					
20150211	7065.09594	3600	53	56	48	48	60	55					
20150212	7066.11174	3600	57	63	54	54	68	65					
20150213	7067.12605	3600	53	57	49	49	62	57					
20150214	7068.12689	3600	58	67	55	55	76	71					
20150215	7069.07940	3600	63	66	54	54	66	61					
20160129	7417.15243	3600	87	84	63	67	89	74	28	17	10	7	6
20160226	7445.11321	3600	84	81	61	60	85	71	29	17	10	6	6
20160227	7446.07349	5400	151	145	110	107	151	125	48	30	16	10	10

Strassmeier et al. (2000) discovered that V1355 Ori exhibited very strong emission over the local continuum in Ca II H&K lines, and showed weak emission in one single H_α spectra. Strassmeier (2000) detected a strong flare in the H_α line during doppler imaging observation in 1998. He also found that the H_α profile varies against the orbital phase during the three observing epochs in 1997, 1998, and 1999 (Strassmeier 2000). The profile of the H_α line appear to be quite complex, ranging from an extremely strong emission due to flaring emission with a central sharp absorption, and to partial emission with a red absorption component, and to very narrow absorption (Strassmeier et al. 2000; strassmeier 2000). No analysis has been carried out regarding the H_δ, H_γ, H_β, Na I D_1, D_2, and Ca II IRT lines to date.

It is of great importance to determine the behaviour of chromospheric active indicators, and chromospheric active properties and evolution (Hall 2008; Montes 2004; Zhang et al. 2015; Zhang et al. 2016a,b; etc). In this paper, we present new high-resolution observations of V1355 Ori. We analysed the properties of chromospheric activity for the Ca II H&K, H_δ, H_γ, H_β, Na I D_1 D_2, H_α and Ca II IRT lines.

2 OBSERVATIONS

We obtained new high-resolution spectra for V1355 Ori in twelve nights between Jan. 22 and Feb. 15, 2015 using the Coude Echelle Spectrograph and a 2048×2048 pixels Tektronix CCD detector of the 1.8 meter telescope (Rao et al. 2008; Wei et al. 2000) at the Lijiang station of Yunnan Observatories. The slit width of the spectrograph on the 1.8-m telescope was 61.7-μ m. The resolution is approximately 50,000 in terms of the FWHM of arc lines. The spectral resolution is approximately 50, 000 and the spectral wavelength region is approximately 5760-11960 $\mathring{A}$ (Zhang et al. 2016a). The limiting magnitude of the equipment is about 11.5 mag at present. In order to obtain synthesized spectra depicting the stellar photospheric contribution of V1355 Ori, we specifically chose and observed four single inactive stars (HR 617 (K1 III), Beta Gem (K0 III), HR 4182 (G9 IV) and HR 495 (K2 IV)) with spectral types and luminosity classes similar to those of V1355 Ori. HR 617 and Beta Gem were also used as the reference stars for V1355 Ori to discuss the chromospheric activity (Strassmeier 2000; etc).

We also observed V1355 Ori using the new high-resolution spectrograph and a 4096×4096 pixels Tektronix CCD detector on the 2.4-m telescope (Fan et al. 2015) at the Lijiang station of Yunnan Observatories in three nights (Jan. 29 and Feb. 26 and 27, 2016). The slit width used in this part of the observation was 62.5 μ m. The spectral wavelength region is approximately 3890-10600 $\mathring{A}$. The spectral resolution in terms of the FWHM of the arc comparison lines is 0.110 $\mathring{A}$ in the Ca II H&K lines, 0.145 $\mathring{A}$ in the H_δ, 0.117 $\mathring{A}$ in the H_γ, 0.152 $\mathring{A}$ in the H_β, 0.146 in several metal lines (6400-6510$\mathring{A}$), 0.147 $\mathring{A}$ in the He I D_3, Na I D_1, D_2 lines, 0.150 $\mathring{A}$ in the H_α line, 0.212 $\mathring{A}$ in the Ca II IRT 8498 $\mathring{A}$ line, 0.245 $\mathring{A}$ in the Ca II IRT 8542 $\mathring{A}$ line, and 0.227 $\mathring{A}$ in the Ca II IRT 8662 $\mathring{A}$ line. The



Fig. 1 All the observed spectra for V1355 Ori using 1.8-m telescope in the He I D_3, Na I D_1 D_2, H_α, and Ca II 8542 lines. All these spectra were vertically shifted by 1.

corresponding spectral resolution is about 43, 000 according to the FWHM of the arc comparison lines. The limiting magnitude of the equipment is about 12 mag at present. We also observed HR 617 (K1 III) as a reference star for V1355 Ori to discuss the chromospheric activity.

We followed the standard process to reduce the spectra using the Image Reduction and Analysis Facility (IRAF) package, which contains tools to perform CCD image trimming, bias subtraction, flat-field correction, removal of cosmic rays, background subtraction, and multi-spectrum extraction. The wavelength was calibrated using the Th-Ar lamp, and the observed spectra were normalized by a low-order polynomial function. The spectra of V1355 Ori from Lijiang 1.8-m telescope are illustrated



(a) Ca II 8662 and 8542 lines



(b) Ca II 8498 and metal lines



(c) H_α and Na I D_1 D_2 lines





(e) Ca II H&K and H_δ lines

Fig. 2 The observed, synthesized, and subtracted spectra of V1355 Ori from the 2.4-m telescope in several spectral lines (Ca II H&K, H_δ, H_γ, H_β, Na I D_1 D_2, H_α and Ca II IRT lines), and metal lines in the spectral wavelength region of 6400-6500 Å. All these spectra were vertically shifted by several units.

in Fig. 1 and the spectra from 2.4-m telescope are plotted in Fig.2. We list our observation log of V1355 Ori in Table 1, which includes observational date, the Heliocentric Julian Date (HJD), exposure times, and signal to noises ratios of the chromospheric active lines.

3 SPECTROSCOPIC ANALYSIS

For the observed spectra (Figs. 1 and 2), all the H_α spectral lines exhibit variable behaviour (filled-in absorption, partial emission with a center absorption component, or emission above continuum). The Na I lines demonstrate deep absorptions and the Ca II IRT lines exhibit filled-in absorptions with minor self-reversal core emissions. The Ca II H&K lines show obvious emissions above continuum. The H_δ, H_γ, and H_β lines exhibit absorptions.

We analysed the spectra of V1355 Ori using a synthetical spectral subtraction technique (Barden 1985; Montes et al. 1995). The synthesized spectra of V1355 Ori were obtained from rotationally broadened and radial-velocity shifted spectra of a single inactive star with similar spectra type and luminosity class as our object. The rotational velocity value (vsini= 39.6 km s^{-1}) of V1355 Ori was determined using some single metallic spectra lines in the wavelength ranges of 6387-6487 Å (Fig. 2). We chose Beta Gem for the 1.8m telescope and HR 617 for the 2.4m telesccope for this purpose. Our result is similar to the previous results of 40 $\pm$ 0.5 km s^{-1} derived by Strassmeier (2000) and 46 km s^{-1} (Osten & Saar 1998; Strassmeier et al. 1999). All the subtracted, observed, and synthesized spectra are shown in Figs. 2 and 3. The Ca II IRT lines have a poorer quality than the Na I and H_α lines because of shorter exposure time and poorer weather condition at the time of observation. For Ca II H&K lines, we did not used spectral subtraction technique to analyse them because their emission over continuum and low S/N ratio. The subtraction spectra of the Na I D_1, D_2 lines show weak emissions, while all subtracted spectra of the H_α line exhibit obvious emission over the continuum. The excess spectra for the Ca II IRT lines display emissions. There are no excess emission in the H_δ, and H_γ lines, but there is excess emission in the H_β line. The EWs were calculated by integrating them above the emission lines using the SPLOT package of IRAF. The methods for calculating the EWs and their uncertainties have been published in our previous papers (Zhang & Gu 2008; Zhang et al. 2014; Zhang et al. 2015; etc). For



Fig. 3 Observed and synthesized (left), and subtracted (right) spectra of V1355 Ori using 1.8-m telescope.

V1355 Ori, the uncertainties of the EWs are underestimated because the S/Ns for these spectra are low and we do not consider the effect of the error of continuum.

The values of the HJD, orbital phase, the excess EWs of chromospheric activity indicators, and the ratios of EW_{8542}/EW_{8498} and E_{H_α}/E_{H_β} are listed in Table 2. The orbital phases were calculated using the function: HJD = 2450540.365 + 3.87192*E (Strassmeier 2000). We also plotted the EWs of V1355 Ori against the HJD or phase in Fig. 4, where different symbols represent different chromospheric active indicators.

4 DISCUSSIONS AND CONCLUSIONS

The spectra we observed show obvious emissions above continuum in the Ca II H&K lines, and variable behaviour for the H_α line. Our results further confirm those behaviour observed in the Ca II H&K lines (Strassmeier et al. 2000) and the H_α line (Strassmeier 2000; Strassmeier et al. 2000). These spectra

Table 2 The values of chromospheric emissions of V1355 Ori

HJD(245,)	Phase	$EW_{Ca_{II}K}(\AA)$	$EW_{Ca_{II}H}(\AA)$	$EW_{Na_ID_1}(\AA)$	$EW_{Na_ID_2}(\AA)$	$EW_{H_\alpha}(\AA)$	$EW_{H_\beta}(\AA)$	E_{H_α}/E_{H_β}	$EW_{Ca_{II}8498}(\AA)$	$EW_{Ca_{II}8542}(\AA)$	$EW_{Ca_{II}8662}(\AA)$	EW_{8542}/EW_{8498}
7045.22327	0.009	-	-	0.093 ± 0.011	0.055 ± 0.010	1.257 ± 0.180	-	-	1.107 ± 0.016	1.037 ± 0.189	1.31 ± 0.057	0.937 ± 0.154
7046.23188	0.269	-	-	0.087 ± 0.002	0.058 ± 0.010	1.200 ± 0.016	-	-	0.464 ± 0.029	0.646 ± 0.076	1.393 ± 0.133	1.392 ± 0.132
7050.17989	0.289	-	-	0.086 ± 0.008	0.049 ± 0.001	1.663 ± 0.043	-	-	0.967 ± 0.070	1.306 ± 0.148	0.846 ± 0.086	1.351 ± 0.113
7058.14409	0.346	-	-	0.115 ± 0.009	0.057 ± 0.009	1.276 ± 0.023	-	-	0.496 ± 0.055	0.810 ± 0.106	0.959 ± 0.11	1.633 ± 0.172
7060.11388	0.855	-	-	0.137 ± 0.011	0.063 ± 0.001	2.012 ± 0.098	-	-	1.089 ± 0.087	1.295 ± 0.470	1.455 ± 0.069	1.189 ± 0.375
7063.11839	0.631	-	-	0.063 ± 0.013	0.011 ± 0.005	1.371 ± 0.034	-	-	0.614 ± 0.009	0.791 ± 0.144	0.891 ± 0.039	1.288 ± 0.226
7064.13065	0.892	-	-	0.061 ± 0.002	0.016 ± 0.012	1.163 ± 0.064	-	-	0.971 ± 0.047	0.890 ± 0.338	0.945 ± 0.259	0.917 ± 0.290
7065.09594	0.141	-	-	0.043 ± 0.003	0.032 ± 0.007	1.249 ± 0.025	-	-	1.003 ± 0.094	1.028 ± 0.431	1.086 ± 0.095	1.025 ± 0.340
7066.11174	0.404	-	-	0.090 ± 0.009	0.056 ± 0.005	1.425 ± 0.038	-	-	1.039 ± 0.015	0.992 ± 0.102	1.131 ± 0.149	0.954 ± 0.082
7067.12605	0.666	-	-	0.082 ± 0.008	0.017 ± 0.001	1.227 ± 0.011	-	-	0.869 ± 0.058	1.183 ± 0.262	1.113 ± 0.003	1.361 ± 0.265
7068.12689	0.924	-	-	0.086 ± 0.008	0.023 ± 0.001	1.561 ± 0.034	-	-	0.896 ± 0.014	1.028 ± 0.555	1.386 ± 0.062	1.147 ± 0.608
7069.07940	0.170	-	-	0.108 ± 0.003	0.073 ± 0.005	1.765 ± 0.003	-	-	0.544 ± 0.090	0.891 ± 0.347	1.099 ± 0.228	1.638 ± 0.576
7417.15243	0.593	3.992 ± 0.279	4.261 ± 0.167	-	-	1.255 ± 0.056	0.273 ± 0.010	5.789 ± 0.046	1.054 ± 0.141	1.619 ± 0.091	1.271 ± 0.252	1.536 ± 0.119
7445.11321	0.288	3.608 ± 0.092	2.620 ± 0.230	-	-	1.232 ± 0.030	0.267 ± 0.007	5.811 ± 0.011	0.605 ± 0.024	0.869 ± 0.093	0.926 ± 0.012	1.436 ± 0.097
7446.07349	0.536	4.214 ± 0.160	3.834 ± 0.091	-	-	1.546 ± 0.100	0.235 ± 0.005	8.285 ± 0.360	0.614 ± 0.018	0.872 ± 0.038	0.960 ± 0.020	1.420 ± 0.020

show deep absorptions in the Na I D_1 D_2 lines, absorptions for the H_δ, H_γ, H_β, and minor self-reversal emissions in the Ca II IRT absorption lines. These features further demonstrate the chromospheric activity of V1355 Ori. The subtracted spectra exhibit no excess emission in the H_δ, and H_γ lines, trace amount of emissions in the Na I lines, excess emission in the H_β line, clear emission in the H_α line, and weak excess emissions in the Ca II IRT lines. The maximum EW of our H_α line is smaller than that of the large H_α flare at HJD 2450909.6 (Strassmeier 2000).

The value of the ratio (EW_{8542}/EW_{8498}) is also an indicator of the chromospheric activity. For V1355 Ori, the value of EW_{8542}/EW_{8498} is in the range of 0.9-1.7. Smaller values indicate optically thick emissions from possible stellar plage event. These values are also consistent with other behaviour of chromospheric activity indicators. Similar values were also discovered in many chromospheric active stars (Cao & Gu 2005; Arévalo & Lázaro 1999; Gu et al. 2002; Montes et al. 2001; Zhang et al. 2014, 2015; Zhang et al. 2016a; etc).

We calculated the ratio of E_{H_α}/E_{H_β} for V1355 Ori in our 2016 run (Table 2). The ratio of excess emission E_{H_α}/E_{H_β} was corrected with the function $\frac{E_{H_\alpha}}{E_{H_\beta}} = \frac{EW_{H_\alpha}}{EW_{H_\beta}} * 0.2444 * 2.512^{(B-R)}$ from Hall & Ramsey (1992). The color index of V1355 Ori (B-R)=1.78 were assumed by its spectral type. Buzasi (1989) obtained that low ratios (1-2) can be achieved both in plages and prominences viewed against the disk, but high values (3-15) can only be achieved in prominence-like structures viewed off the stellar limb. We obtained a mean value of 6.628 ± 0.139 for E_{H_α}/E_{H_β} for V1355 Ori. The ratio E_{H_α}/E_{H_β} ($\gtrapprox 3$) implies that the emission of the Balmer lines originates from prominence-like material (Buzasi 1989; Hall & Ramsey 1992).

It is unpractical for us to carry out a detailed study on the chromospheric rotational modulation of V1355 Ori because of data acquisition limitations in the night (we normally only have a partial night) and the total time span required. We had twenty two nights of the 1.8-m telescope for studying V1355 Ori, which corresponds to approximately six orbital periods. Our results do not show any obvious phase modulations in all the chromospheric active indicators lines (Fig. 4). Chromospheric intensity as well as the presence of circumstellar material may dilute or even suppress the rotationally modulated plage signature (Strassmeier 2000). As illustrated in Fig. 4, there is a weak time-variation of excess emission EWs in the Na I D_1 D_2 Ca II IRT, and H_α lines (which is the strongest). The increase of EWs is marked by red arrows in Fig.4. All these indicators are consistent with each other. These phenomena can be explained by plage events, which is consistent with the observed behaviour of these chromospheric active indicators. In the future, further high-resolution spectra will be required to study the chromospheric activity cycle of V1355 Ori, such as those of photospheric cycle reported by Savanov & Strassmeier 2008.



Fig. 4 EW light curves of chromospheric excess emissions of V1355 Ori vs. HJD (Left) and the orbital phases (right). Red arrows mark the events where EWs increased significantly.

Acknowledgements This work was supported by the Astronomic Joint Fund of NSFC and CAS (U1431114 and 11263001), and science and technology innovation team of Guizhou province (Nos. 20154017). This work was also supported by the Natural Science Foundation of the GuiZhou Province office of Education (Grant No. 2014298). We acknowledge the support of the staff of the Lijiang 2.4m telescope. Funding for the telescope was provided by CAS and the People's Government of Yunnan Province.

References

Arévalo, M. J., & Lázaro, C. 1999, AJ, 118, 1015
Barden, S. C. 1985, ApJ, 295, 162
Buzasi, D. L. 1989, PhD thesis, Pennsylvania State Univ.
Cutispoto, G., Pallavicini, R., Kuerster, M., et al. 1995, A&A, 297, 764
Cao D. T., Gu, S. H., 2015, MNRAS, 449, 1380C
Eker, Z., Ak, N., Filiz, B., et al. 2008, MNRAS, 389, 1722
Fan,Y.F. et al. 2015, RAA, 15,918F
Gu, S. H., Tan, H. S., Shan, H. G., et al. 2002, A&A, 388, 889
Hall, J. C., & Ramsey L. W. 1992, AJ, 104, 1942
Hall, J. C. 2008, Living Rev. Solar Phys., 5, 2
Montes, D., Crespo-chaón I., Gálvez, M. C. et al. 2004, LNEA, 1, 119
Montes, D., Fernández-Figueroa, M. J., de Castro, E., et al. 1995, A&AS, 109, 135
Montes, D., López-Santiago, J., Gálvez, M. C., et al. 2001, MNRAS, 328, 45
Osten, R. A., & Saar, S. H. 1998, MNRAS, 295, 257
Pounds, K. A., Allan, D. J., Barber, C., et al. 1993, MNRAS, 260, 77
Rao, C. H., Jiang, W. H., Zhang Y. D., et al. 2008, SPIE, 7015E, 5YR
Strassmeier, K. G., Serkowitsch, E., Granzer, Th., et al. 1999, A&AS, 140, 29
Strassmeier, K. G., Washuettl, A., Granzer, Th., et al. 2000, A&AS, 142, 275
Strassmeier, K. G. 2000, A&A, 357, 608
Savanov, I. S., & Strassmeier, K. G. 2008, AN, 329, 364
Voges, W., Aschenbach, B., Boller, Th., et al. 1999, A&A, 349, 389
Wei, K., Zhang, X. J., Xian, H., et al. 2000, Chinese Optics Letter, 8, 1019
Zhang, L. Y., & Gu, S. H. 2008, A&A, 487, 709

Zhang, L. Y., Pi, Q. F., Zhu, Z. Z., et al. 2014, NewA, 32, 1
Zhang, L. Y., Pi, Q. F., Zhu, Z. Z. 2015, RAA, 15, 252
Zhang, L. Y., Pi, Q. F., Han, Xianming L. et al. 2016a, MNRAS, 459, 854
Zhang, L. Y., Pi, Q. F., Han, Xianming L. et al. 2016b, NewA, 44, 66
Zirin, H. 1988, Astrophysics of the Sun, Cambridge University Press